# Assessment of Students' Knowledge about Environmental Health in College of Health and Medical Technology: A Cross-Sectional study

تقييم معارف الطلبة حول صحة البيئة في كلية التقنيات الصحية والطبية : دراسة مقطعية

Sulaf A. Hussein, MsC\* Suhair M. Hasoon, M, Tec\*\*

#### المستخلص:

الهدف: الهدف من الدراسة هو تحديد مستوى معرفة الطلبة حول صحة البيئة.

المنهجية: دراسة مقطعية أجريت في كلية التقنيات الصحية والطبية في مدينة بغداد خلال الفترة من امارس ولغاية ايوليو ١٠١٧وقد تم جمع البيانات عن طريق التسجيل الذاتي للاستبيان الذي صمم مسبقا للحصول على المعلومات الاجتماعية والديمغرافية مثل (العمر،الجنس، القسم، سنة الدراسة). النتائج: أعلى نسبة للطلبة كانوا في المرحلة الثانية و يليها المرحلة الثائة وكانت اعلى معدلات الطلبة هم الذين لديهم اقل مستوى معرفي ويتبع ب المستوى المعرفي المعرفي المعرفي المعرفي العاليافتيات هم اعلى مستوى معرفي مقارنة مع الذكور الذين هم مستواهم المعرفي متوسط. الطلبة من قسم التحليلات المرضية بلغوا اعلى مستوى معرفي يليهم طلبة قسم التخدير واقل مستوى معرفي كانت في قسم العلاج الطبيعي . وجود فرق معنوي ما بين انخفاض مستوى المعرفة في السنة الأولى (٧,٧٠٪) والسنة الثانية (٧,٠٥٪)، بينما كانت اعلى نسب للمعرفة بشكل رئيسي لطلبة المرحلة الرابعة (٩,٨٪) و متوسط مستوى للمعرفة في السنة الثالثة حيث بلغت نسبتها (٣,٣٥٪) تليها السنة الرابعة بنسبة ٥٠٪. التوصيات: كل اقسام الكلية يجب ان يدرس فيها مادة صحة البيئة لما لها من اهمية وليس فقط قصرا على طلبة قسم صحة المجتمع.

## **Abstract:**

Objective: the aim of this study is to determine the level of students' knowledge about the environmental health.

**Methodology:** The cross-sectional study was conducted at the College of Health and Medical Technology in Baghdad city during the period from 1st march till 1st of July 2012. Data was collected by self-recording of a previously designed questionnaire to obtain socio-demographic information such as (age, gender, department, year of grade).

**Results:** The highest rate of students were in the 2nd year followed by the 3rd year, highest rate of students had low level of knowledge followed by intermediate level of knowledge, while lowest rate of students on had high level of knowledge .Females had higher level of knowledge compared to males who had intermediate Level of knowledge; Students of pathological analysis department had high level of knowledge among other students, followed by students of anesthesia department. The lowest rate was among department the physiotherapy. A significant association between that low level of knowledge was among 1st (57.7%) and 2nd (50.7%) year students mainly , while high level of knowledge was mainly among 4th year (9.8%), and the intermediate level of knowledge was among 3rd year students in a rate of (53.3%) and 4th year in a rate of 50%.

**Recommendations:** Include all the department of the college in environment health lectures as the issue is important for all student and not only the community health students.

Key words: Assessment, Students, Environmental health, Knowledge.

<sup>\*</sup>Assistant Professor, Community Health Department, College of Health and Medical Technology

<sup>\*\*</sup>Instructor, Community Health Department, College of Health and Medical Technology

## Introduction:

nvironmental Education (EE) has been defined by the United Nations Educational, and Cultural Scientific Organization (UNESCO) as "a learning process that increases people's knowledge and awareness about the environment and associated challenges, develops the necessary skills and expertise to address the challenges, and fosters attitudes, motivations, and commitments to make informed decisions and take responsible action" (1). The Decade of Education for Sustainable Development (DESD, 2005-2014) was proclaimed by the General Assembly of the United Nations in December 2002 in resolution 57/254. The basic vision of the Decade is a world where everyone has the opportunity to benefit from education and learn the values, behaviors and lifestyles required for a sustainable future. Education for sustainable development addresses the issues of natural resources (water, energy, agriculture, housing, biodiversity, etc) and develops the understanding of the interdependence and fragility of the Earth systems. This will enable learners to adopt new behaviors in the protection and use of natural resources, which are essential for human development and survival (2). In recent years, environmental sensitivity has been centered in the of domain environmental education environment specialists and it has been used within a sense of emphatic perspective into the environment (3). A large topic of interest within the environmental education realm is establishing how personal traits or lifestyle factors contribute to a person's environmental attitudes and behaviors. A general attitude can be defined as something which "must be perceived by the individual as connected in some meaningful way to a specific situation to serve as a basis for an evaluative reaction in that situation (4). Knowledge is commonly seen as a necessary precondition for a person's behavior. Consistent with this, most educational interventions rely on knowledge

transfer. However, for the most efficient informational strategies for education, it is essential that we identify the types of knowledge that promote behavior effectively and investigate their structure (5). Student's environmental awareness is one of the most important indicators for displaying national civilization. It reflects many aspects of environmental status, such as personal consideration and behavior, public capacity, and the local citizens, attitude towards sustainable society as a whole, (6). An understanding of the nature of environmental problems thereby is fundamental for any approach to addressing their solutions and this necessitates the documentation of our students' present level of environmental consciousness and understanding of related concepts (7). Thus, this paper discusses the findings of a survey on university students' level of environmental attitudes, behaviors and knowledge and its implications for education in the light of sustainable development (7), the aim of this study to identify environment health knowledge of student.

# Methodology:

The study design was cross - sectional one, conducted in College of health and medical technology, for the period from 1<sup>st</sup> march till 1<sup>st</sup> of July 2012. The sampling method was stratified random sampling. And thus sample size was (807). Data collection was by self-recording of a previously designed questionnaire that consisted of several Components, namely demographic background and Knowledge; which tested the knowledge of the students regarding some environmental topics such as definition of the environment, global environmental problems, water pollution, air pollution, solid waste and energy. These topics were tested through 25 multiple choice questions, the source and Frequency of exposure of students to environmental information outside school or college.

Each question was allowing for one correct answer only. A score was calculated for every student. Scores were found to range between 8/25 and 22/25. A score of  $\leq$  69% was considered poor, 70-80% was considered moderate and a score of 81% or more was considered good. Then rate of poor and moderate and high percentages were calculated and students' knowledge was classified accordingly.

Data were analyzed using descriptive statistics (frequencies and percentages), and analytic statistics (chi-square test for association between two variables with results being considered as statistically significant when the p value was < 0.05.

## **Results:**

Table 1. Distribution of Students' Characteristics (No. 807)

Characteristic of sample	Frequency	Percentage
Gender		
Male	360	44.6
Female	447	55.4
Grades		
1st year	182	22.6
2nd year	219	27.1
3rd year	212	26.3
4th year	194	24
Department		
Technical Pathological analysis	161	20
Technical Dentist	74	9.2
Technical x- ray	134	16.6
Technical Ophthalmology	96	11.9
Technical Anesthesia	108	13.4
Technical Physiotherapy	81	10
Technical Community health	153	19

No: sample size

Females constituted 55.4% of the sample as shown in table 1, the highest rate of students were in the 2nd year (27.1%) followed by the 3<sup>rd</sup> year in a rate of (26.3%). 20% of students were from the department of technical pathological analysis and 19% from the technical community health department, the lowest rate was the technical dentist department (9.2%).

Table 2. Distribution of Students According to their Knowledge

Level of knowlege	Frequency	Percent
Poor	395	49
Fair	372	46
Good	40	5
Total	807	100

Table 2 shows that the highest rate of students had low level of knowledge (49%) followed by a rate of 46% for intermediate level of knowledge and only 5% of students on had high level of knowledge.




Figure 1. Level of Knowledge for Students

Table 3. Rate of Respondents with Correct Answer for Knowledge of Certain Issues on Environmental Health

Type of knowledge question	Rate of correct answers
Natural environment	90.4
Environmental pollutants	52.4
Greenhouse gases	17.7
Ozone depletion	54.2
Desertification	4.7
Solid waste disposal	40.1
Air pollutant	85.4
Global warming	39.14
Loss of natural Resources	27.15
Recycling of waste material	37.7

Table 3 shows that students had high score of knowledge about questions related to natural environment (90. 4%) followed by knowledge about air pollutants (85.4%) while only 4.7% knew what desertification means.

Table 4. Relationship between Students' Knowledge and Gender

Candan		Level of knowledg	e	Total	Duglug
Gender	Poor	Fair	Good	Total	P value
Male	184	160	16	360	
	51.1%	44.4%	4.4%	100%	χ²= 2.043
Female	206	212	24	447	p <u>&lt;</u> 0.509
	46.1%	47.4%	5.4%	100%	N.S
Total	383	372	40	807	
	47.5%	46.1%	5%	100%	

%= percent, P= probability level, N.S= No significant

Females had higher level of knowledge as shown in table 4 as the rate for intermediate and high level was 47.4% and 5.4% respectively as compared to males who had a rate of 44.4% for intermediate Level of knowledge and a rate of 4.4% for high level. The result was statistically not significant as P value was more than 0.05.

Table 5. Relationship between Students' Knowledge and Class

Class	L	Level of knowledge			
	Poor	Fair	Good		
1 <sup>st</sup>	105	66	1	182	
	57.7%	36.3%	.5%	100%	
2 <sup>nd</sup>	111	96	12	219	
	50.7%	43.8%	5.5%	100%	χ²= 57.79
3 <sup>rd</sup>	90	113	8	212	p <u>&lt;</u> 0.000
	42.5%	53.3%	3.8%	100%	H.S
4 <sup>th</sup>	77	97	19	194	
	39.7%	50%	9.8%	100%	
Total	383	372	40	807	
	47.5%	46.1%	5%	100%	

%= percent, χ²= chi-squared test, P= probability level, H.S= highly significant

Table 5 shows that low level of knowledge was among  $1^{st}$  (57.7%) and  $2^{nd}$  (50.7%) year students mainly while high level of knowledge was mainly among  $4^{th}$  year (9.8%), and the intermediate level of knowledge was among  $3^{rd}$  year students in a rate of 53.3%) and  $4^{th}$  year in a rate of 50%. The result is statistically highly significant as p value was less than 0.05.

**Table 6.** Relationship between of Students' Knowledge and Type of Study

Department	Level of knowledge			Total	p- value
	Poor	Fair	Good		
Technical pathological analysis	88	48	25	161	
	54.7%	29.8%	15.5 %	100%	
Technical dentist	51	22	1	47	
	68.9%	29.7%	1.4%	100%	χ²= 149.1
Technical x- ray	59	78	2	134	p <u>&lt;</u> 0.000
	61.5%	58.2%	1.5%	100%	H.S
Technical ophthalmology	59	36	1	96	
	61.5%	37.5%	1%	100%	
Technical Anesthesia	38	65	5	108	
	35.2%	60.2%	4.6%	100%	

**Table 6. Continues** 

Technical physiotherapy	56	25	0	81
	69.1%	30.9%		100%
Technical community health	49	98	6	153
	32%	64.1%	3.9%	100%
Total	395	372	40	807
	48.9%	46.5%	5%	100%

%= percent, χ²= chi-squared test, P= probability level, H.S= highly significant

Table 6 shows that students of technical pathological analysis department had high level of knowledge in a rate of 15.5% which is the highest among other students, followed by students of technical anesthesia in a rate of 4.5%. The lowest rate was among department of technical physiotherapy 0% .The result is statistically highly significant as p value was less than 0.05.

## **Discussion:**

The characteristic of sample reflects distribution of students in the college of health and medical technology according departments as the technical laboratory analysis department is the largest in terms of number of students followed by the technical community health department. Our current study, the students' written responses to closed ended questions revealed varying degrees of knowledge concerning environmental health issues. In terms of students' baseline knowledge, results showed that in general, students have more knowledge about more general older environmental topics (Environmental Issues) than the specific topics or terminologies, as for constituent of natural environment 90.4% answered correctly, also causes of air pollution was known to 85.4% of students, The results were consistent with the findings of a world-wide study which showed that the level of knowledge on environmental issues among surveyed undergraduates was not satisfactory and there was a large knowledge gap on almost every aspect from environmental tools and technologies (7) and 78.3% of Italian population were aware that outdoor air pollution was associated with increase daily mortality from respiratory diseases (8). The greenhouse effect that is produced by industry and agriculture and is believed to have a major role in global

warming<sup>(9)</sup>was known to 17.7% of students in the current study only while students of Malaysia had higher rate of knowledge about this issue amounting to 82% (10) and in Italy the rate was 78.9%<sup>(8)</sup>. The causes of ozone depletion were known to 54.2% of students in the study sample, which is similar to results of a study in China that 55.8% of students knew chlorofluorocarbon emissions from refrigerators was the main cause for ozone depletion (11) and so was most first year medical students in Ankara<sup>(12)</sup>. Recent issue like global warming was known for only 39.14% of students and desertification which is an important problem nowadays is known for only 4.7% of students a rate close to that of medical students in Ankara (11). Comparing these results with those of a study in America, it was found that 14% of their population were alarmed about the effect of global warming and 31% were every concerned that greenhouse gases in the affect atmosphere the average global temperature (13) and the Malaysian population recognized the fact that deforestation will cause biological imbalance in a rate of 53%<sup>(14)</sup>.

Management of solid waste reduces or eliminates adverse impacts on the environment and human health and supports economic development and improved quality of life. Knowledge about solid waste and its management was known to 40.1% of students in

the current study, this result was similar to that of a study in Iran among students of medical Sciences whose rate of knowledge about this issue was 44% (15) . Resource recovery offers alternative to waste management but recycling was known to only 37.7% of them, the result is low compared to that of poor people in Malaysia who had an adequate knowledge about waste management and recycling (16) .Natural resources are derived from the environment. Some of them are essential for our survival while most are used for satisfying our wants (17). Knowledge of students about natural resources conservation was very poor (27.15%) compared to Omani's knowledge about this issue which was in a rate of (62.5%) (18). Results of the current study revealed a non-significant difference in gender distribution regarding level of knowledge although there was higher level of knowledge among females than males. In Iran there was higher knowledge of males over females (15). Studies in China had demonstrated that women had participation in environmental behaviors inside of the home (e.g., recycling), while outside of the home (e.g., environmental organization donations) no gendered patterns were exhibited (11). Years of study had effect on the level of knowledge of students as 4th year students had higher rate but the difference is not great as one should expect from students in the health field and this is further emphasized by the higher results of the Technical pathological analysis department students over the technical community health students who should have the highest score according to the subjects in their curriculum, but the results were opposite to expectations indicating an error either in the topics of environmental health lessons or the perception of students to these topics.

## **Recommendation:**

- 1. Development and implementation of environmental education programs as part of the regular school curriculum
- 2. Environmental education should be part of the curriculum of all departments of the college to ensure the consistency of environmental practices among students, and it should address both local environmental issues and global environmental problems.
- 3. Pamphlets and guides distributed to the students may help to remind them about the importance of healthy environment and how to achieve that.
- 4. Audits biannually can press students to improve performance. This action needs technical support provided by the college to launch a sustainability audit project involving training and a manual.

### References:

- US Environmental Protection Agency. Environmental education background and history. [Cited 2009, Feb 20th]. Available from: http://www.epa.gov/enviroed/eedefined.html
- Bory-Adams A. UNESCO's role, vision and challenges for the UN decade of education for sustainable development (2005-2014).
   Connect; 31 (1-2): 1-5, 2006.
- 3. Kim, K. O. "An inventory for assessing environmental education curricula". *The Journal of Environmental Education*, 34 (2): 12–18, 2003.
- Marczyck, G. R., DeMatteo, D., & Festinger, D.
   Essentials of research design and methodology. Book, John Wiley & Sons; Hoboken, NJ, 290, (2005).
- Frick J; Kaiser FG; Mark W. Environmental knowledge and Conservation Behavior: Exploring Prevalence and Structure in a Representative Sample. Personality and Individual Differences J. 37(8): 1597-61, 2004

- Kaiser FG. Ecological behaviors, s Dependency on Different form of Knowledge. Applied psychology: An International Review 52: 598-613, 2003.
- Azapagic, A., Perdan, S., & Shallcross, D. How much do engineering students know about sustainable development? The findings of an international survey and possible implications for the engineering curriculum. European Journal of Engineering Education, 30, 1–19, 2005.
- 8. Bianco A; Nobile C; Gruisci F; Pavia M. knowledge and perceptions of the health effects of encironmental hazards in the general population in Italy. International J. of Hygiene and Environmental Health, 211, issues 3-4: 412-419, 2008.
- Wallington TJ; Ole SJ;,Ellie J. Highwood JN. GREENHOUSE GASES AND GLOBAL WARM-ING, in Environmental and Ecological Chemistry, [Ed. Aleksandar Sabljic], in Encyclopedia of Life Support Systems (EOLSS), Developed under the Auspices of the UNESCO, Eolss Publishers, Oxford ,UK, (2004), [http://www.eolss.net]
- 10.Syde Idros SN. Exploring environmental behaviors, attitudes and knowledge among university students: positioning the concept of sustainable development within Malaysian education.
- 11.Xiao C, Hong D. gender differences in environmental behaviors in China. *Popul Environ* 32:88–104, 2010.
- 12.Guler C; Altintash H; Temel F; Ahrabi AF, et al. Evaluation of the environmental consciousness of the students in a medical faculty in Ankara. *TAF preventive medicine Bulletin*, 5(4) 151-171, 2005.
- 13.Leiserowitz A and Smith N. knowledge of climate change across global warming's six Americas. Yale University, New Haven. CT:

- Yale project on climate change communication, 2010.
- 14.Haron SA; Paim L; and Yahaya N. Towards sustainable consumption: an examination of environmental knowledge among Malaysians. International Journal of Consumer Studies, 29, 5, September 2005, pp426–436
- 15.Ehrampoush, and Bagiani Moghadam. Survey of knowledge,attitude and practice of Yazd University of Medical Science students about Solid waste disposal and recycling. Iranian J of Env. Health Science & Engineering, 2, issue: 2, pp26-30, 2005.
- 16. Murad W; Siwar Ch. Knowledge, attitude and behavior of the Urban Poor Concerning Solid Waste Management. *J of Applied Sciences*, 7 (22): 3356-3367, 2007.
- 17. Schilling M and Chiang L. The effect of natural resources on sustainable development policy:
  The approach of non-sustainable externalities. Energy Policy, 39: 990-998, 2011.
- 18.Sabah A. Abdul-Wahab. A Preliminary Investigation into the Environmental Awareness of the Omani Public and their Willingness to Protect the Environment. American Journal of Environmental Sciences, 4 (1): 39-49, 2008.